arXiv:hep-lat/9909157v3 8 Jan 2000

Factorization of Fermion Doubles on the Lattice.

Pedro J. de A. Bicudo
Departamento de Física and Centro de Física das Interacções Fundamentais, Edifício Ciência,
Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal
email: fbicudo@alfa.ist.utl.pt

We address the problem of the fermion species doubling on the Lattice. Our strategy is to factorize the fermion doubles from the action. The mass term of the Dirac-Wilson action is changed. In this case the extra roots which appear in the action of free fermions in the moment representation are independent of the mass and can be factorized from the fermion propagator. However the gauge couplings suffer from the pathological ghost poles which are common to non-local actions. This action can be used to find a solution of the Ginsparg Wilson relation, which is cured from the non-local pathology. Finally we compare this factorized action with solutions of The Ginsparg Wilson relation. We find that the present is equivalent to the Zenkin action, and that is not quite as local as the Neuberger action.

11.15.Ha

I. INTRODUCTION

Fermion doubles appear in the lattice whenever one calculates the first derivative with the finite difference method and Fourier transforms it. This is an old problem [1], which afflicts lattice simulations of any quantum field theory with light Dirac Fermions. Bosons are healthy because the second derivative has no doubling problem. This problem can be addressed in a simple 1 dimensional lattice. In this case, with a lattice spacing a, the first derivative of a fermion field ψ, is calculated in the first approximation of the finite difference method,

$$\partial\psi_n = \frac{\psi_{n+1} - \psi_{n-1}}{2a} \tag{1}$$

The Fourier transform of an infinite and periodic lattice is the Brillouin zone, a continuous interval $]-\pi, \pi]$, and the functions are transformed as,

$$\psi_n = \int_{-\pi}^{\pi} \frac{dk}{2\pi} e^{in.k}\, \widetilde{\psi}(k) \quad , \quad \widetilde{\psi}(k) = \sum_n e^{-ik.n}\, \psi_n \tag{2}$$

which we substitute in eqs. (1) and find for the derivative,

$$\begin{aligned}\partial\psi_n &= \int_{-\pi}^{\pi} \frac{dk}{2\pi} \frac{e^{+ik} - e^{-ik}}{2a} e^{in.k}\, \widetilde{\psi}(k) \\ &= \int_{-\pi}^{\pi} \frac{dk}{2\pi} i \frac{\sin k}{a} e^{in.k}\, \widetilde{\psi}(k)\end{aligned} \tag{3}$$

The continuum limit is reached when a vanishes. In this case it is natural to replace $n \to x/a$ and $k \to ap$, and we get the correct behavior,

$$\frac{\sin k}{a} \to p \tag{4}$$

However, one should check all the points where the numerator $\sin k$ vanish in the Brillouin zone. In Fig. 1 we see in fact that $\sin k$ has two similar roots in the Brillouin zone, at $k = 0$ and at $k = \pi$. This is the source of the doubler problem.



FIG. 1. In the Brillouin zone, where k is dimensionless, we show the functions $\sin k$, $2\sin\frac{k}{2}$ and $2\tan\frac{k}{2}$.

In the literature many methods to avoid the doubling of fermion species on the lattice are presented. A simple solution of this lattice problem would be convenient for the study of chiral symmetry breaking in QCD and for the study of neutrinos in the Standard Model. Wilson [1] included in the action extra hopping terms which projected out the doublers from the onset. However this was not chiral invariant, and Kogut [2] proposed the staggered fermions to improve chiral invariance. Nielsen and Ninomiya [3] produced a topological no-go theorem concerning chiral invariant lattice models with local charge conservation and a local action. To avoid these restrictions, there are some very interesting solutions in the literature which depart from the simple Wilson action. Kaplan [4] extends the lattice with an extra dimension. and places the physical fermions in a domain wall. 't

Hooft [5] recommends to have continuum fermions with lattice bosons. Neuberger [6] redefines the matrix which is used in the fermion determinant, and finds a topologically nontrivial solution to the Ginsparg-Wilson [7] relation. This opens a new perspective to the study of chiral symmetry [8,9], and the study of the chiral anomaly on the lattice. The method presented in this paper belongs to the class of non-local actions [10–13].

We now show a 1 dimensional example where the doubles are factorized from the propagator. We find it more intuitive to place the fermion fields at the vertices of the lattice zones (dual lattice) because we are tempted to replace the periodic function $\sin k$ by the anti-periodic $2\sin\frac{k}{2}$ which has a single root. To get this new function, which is wider in the momentum representation, see Fig. 1, one has to narrow the spacing $2a$ used in the denominator of the derivative. We will later shift our lattice back to the conventional one, the use of the dual lattice here is not crucial. The first approximation of the finite difference method is now,

$$\begin{aligned}\partial\psi_n &= \frac{\psi_{n+\frac{1}{2}} - \psi_{n-\frac{1}{2}}}{a} \\ &= \int_{-\pi}^{\pi} \frac{dk}{2\pi} i \frac{2\sin\frac{k}{2}}{a} e^{in.k}\ \tilde{\psi}(k)\end{aligned} \tag{5}$$

and we obtain the desired function. We will use the notation $\widehat{\psi}_n$ for the interpolated spinor at the site n of the lattice, which is simply the mean value of the spinors which are placed at the vertices of the zone,

$$\begin{aligned}\widehat{\psi}_n &= \frac{\psi_{n+\frac{1}{2}} + \psi_{n-\frac{1}{2}}}{2} \\ &= \int_{-\pi}^{\pi} \frac{dk}{2\pi} \cos\frac{k}{2} e^{in.k}\ \tilde{\psi}(k)\end{aligned} \tag{6}$$

We find that the constant term (6), in sense that it is not the first derivative term, has a root at $k = \pi$. This is the extra root which was present in the first definition (1) of the derivative.

With the interpolated fields (5) and (6), we define a free Dirac action in 1 dimension,

$$\begin{aligned}S &= a\sum_n\ m\bar{\widehat{\psi}}_n\ \widehat{\psi}_n + \bar{\widehat{\psi}}_n \gamma\partial\psi_n \\ &= a\int_{-\pi}^{\pi} \frac{dk}{2\pi} \bar{\tilde{\psi}}(k) \cos\frac{k}{2}\Big(m\cos\frac{k}{2} + i\gamma\frac{2}{a}\sin\frac{k}{2}\Big)\tilde{\psi}(k)\end{aligned} \tag{7}$$

were the δ relations,

$$\begin{aligned}\int_{-\pi}^{\pi} \frac{dk}{2\pi} e^{ik.(n-n')} &= \delta_{n,n'} \\ \sum_n e^{-in.(k-k')} &= \sum_m 2\pi\delta(k-k'+2\pi m)\end{aligned} \tag{8}$$

were used. We can check that both the mass term (one double root) and the derivative term (two different roots) of the Dirac action (7) have two roots. Thus this action complies with the Nielsen and Ninomiya topological theorem [3]. We see that there are still 2 roots , $\frac{2}{a}\tan\frac{k}{2} = m$ and $k = \pi$ in the Lagrangian (7), but only the first root is dynamical in the sense that it depends on the fermion mass. Nevertheless it is possible to define a propagator G which only shows the correct pole in the denominator, and no double,

$$\begin{aligned}G_{nn'} &\equiv \langle\widehat{\psi}_n\bar{\widehat{\psi}}_{n'}\rangle \\ &= \int_{-\pi}^{\pi} \frac{dk}{2\pi a}\ \frac{e^{ik.(n-n')}}{m + i\gamma\frac{2}{a}\tan\frac{k}{2}}\end{aligned} \tag{9}$$

where the $\cos^2\frac{k}{2}$ in the numerator, which cancels the extra root in the denominator, comes from the interpolation of the fermion fields in the definition of the propagator. The extra root is replaced by a pole. The propagator (9) has the right continuum limit,

$$i\gamma\frac{2}{a}\tan\frac{k}{2} + m \to i\ \not{p} + m. \tag{10}$$

For a finite spacing a, we can see in Fig. 1 that this propagator also has the peculiar property $2\tan\frac{k}{2} > k$, which increases the effect of the ultraviolet cutoff $\frac{\pi}{a}$ in fermions. This is in contradistinction with $\sin k < k$.

In the next Section II we define the factorization procedure in detail, for a Lattice in d dimensions. In section III gauge invariance is implemented. In section IV the factorization is adapted to the Ginsparg-Wilson relation. The conclusion is in Section V. In Appendix A an equivalent factorization is studied, and in Appendix B we study lattices where fermions and antifermions are in separate sites.

II. FACTORIZATION OF THE DOUBLES IN D DIMENSIONS

We now extend the action (7) to the d dimensional case, and factorize the doubles from the propagator and the fermion determinant. In this section we study free fermions which can be exactly solved in the momentum representation. However the factorizing procedure must be applicable in position space, in order to include eventually the gauge fields, and to compute the propagator and the determinant with the standard lattice techniques. At this point we have to leave the dual lattice which was used just for the introduction. In order to write the interpolated spinors, it is convenient to define the class of vectors $\vec{\mu}$, with components $\mu_l = +1\,,\ -1$. In d dimensions, the interpolated spinors are now extended to,

$$\widehat{\psi}_{\vec{n}} = \frac{1}{2^d}\sum_{\mu_l}\psi_{\vec{n}+\frac{1}{2}\vec{\mu}}\ ,\quad \partial_j\psi_{\vec{n}} = \frac{1}{a\ 2^{d-1}}\sum_{\mu_l}\mu_j\psi_{\vec{n}+\frac{1}{2}\vec{\mu}} \tag{11}$$

To translate the fields back to the standard positions $\vec{n}$ we apply a shift of $-\vec{\mu}'/2$ on the action (7). This shift leaves the momentum representation of the action invariant. The action is now,

$$S = \left(\frac{a}{4}\right)^d \sum_{\vec{n},\vec{\mu},\vec{\mu}'} \bar{\psi}_{\vec{n}} \Big(m + \frac{2}{a}\sum_j \mu'_j \gamma_j\Big)\psi_{\vec{n}+\frac{\vec{\mu}'-\vec{\mu}}{2}} \tag{12}$$

where we use the convention of hermitean Dirac matrices γ_i. The action (12) can be written in a matricial form,

$$S = a^d \sum_{\vec{n},\vec{n}'} m\bar{\psi}_{\vec{n}} I_{\vec{n},\vec{n}'}\psi_{\vec{n}'} + a^d \sum_{j,\vec{n},\vec{n}'} \bar{\psi}_{\vec{n}}\gamma_j \partial_{j\,\vec{n},\vec{n}'}\psi_{\vec{n}'} \tag{13}$$

where the sites are labeled by the $\vec{n},\vec{n}'$, which are d-dimensional vectors with integer components. Like the Wilson fermion action [1], this action includes extra hopping terms. In the present case the chiral non-invariant terms are all included in the mass term. In eq. (13) the matrices I and D_j are translational invariant, in the sense that the matrix elements only depend on the distance to the diagonal. The Fourier transform of these matrices depends on a single momentum,

$$I_{\vec{n},\vec{n}'} = \int_{-\pi}^{\pi} \frac{d^d k}{(2\pi)^d} e^{i(\vec{n}'-\vec{n}).\vec{k}}\; \widetilde{I}(\vec{k}) \ , \tag{14}$$

and from eq. (11) we get the Fourier transforms,

$$\widetilde{\partial}_j(\vec{k}) = i\frac{2}{a}\tan\frac{k_j}{2}\widetilde{I}(\vec{k}) \ , \quad \widetilde{I}(\vec{k}) = \prod_l \cos^2\frac{k_l}{2} \ . \tag{15}$$

We define the quark propagator G with the Green equation,

$$\Big(mI + \sum_j \gamma_j \partial_j\Big) G = I \ . \tag{16}$$

This Green equation (16) is an extended version of the normal one. Here we replace the usual identity matrix in the right hand side by the matrix I. The action with no doubles is finally defined,

$$\begin{aligned} S &= a^d \sum_{\vec{n},\vec{n}'} \bar{\psi}_{\vec{n}} D_{\vec{n},\vec{n}'}\psi_{\vec{n}'} \ , \\ D &= G^{-1} = m + \sum_j \gamma_j\, I^{-1}\partial_j \end{aligned} \tag{17}$$

where the matrix M, in the limit of massless quarks is chiral invariant. Like the Neuberger matrix [6], the final matrix D is redefined from the initial action (12). Using the delta relations (8) we find the momentum representation of the action,

$$\begin{aligned} S &= a^d \int_{-\pi}^{\pi} \frac{d^d k}{(2\pi)^d} \widetilde{\bar{\psi}}(k) \widetilde{D}(k) \widetilde{\psi}(k) \ , \\ \widetilde{D}(\vec{k}) &= \widetilde{I}^{-1}(\vec{k}) \Big[m\widetilde{I}(\vec{k}) + \sum_j \gamma_j \widetilde{\partial}_j(\vec{k}) \Big] \\ &= m + i\frac{2}{a}\sum_j \gamma_j \tan\frac{k_j}{2} \ . \end{aligned} \tag{18}$$

We see that the new action has no extra roots and has the correct continuum limit. This result can be compared with the literature. Except for the mass term, we recover the Zenkin action [13], which was derived in the formalism of a path integral with the Weyl quantization.

III. COUPLING TO THE GAUGE FIELDS

We now include the coupling of the fermions to the gauge fields, using the Wilson [1] prescription. In order to have a gauge invariant action, the fermion operators $\bar{\psi}$ and ψ are usually linked by gauge transformation paths when they are in different sites. In the case of a elementary link $\vec{j}$,

$$U_{\vec{n},\vec{n}+\vec{j}} = e^{i\vec{B}(\vec{n}+\vec{j}/2).\vec{j}} \ , \ \vec{B} = \frac{1}{2} a\, g \sum_b \lambda_b . \vec{A}_b \tag{19}$$

where λ_b are the generators of the gauge group, and the gauge-invariant version of $\bar{\psi}_{\vec{n}}\psi_{\vec{n}+\vec{j}}$ is,

$$\bar{\psi}_{\vec{n}} e^{i\vec{B}.\vec{j}} \psi_{\vec{n}+\vec{j}} . \tag{20}$$

In the present case we have to connect $\bar{\psi}_{\vec{n}}$ and $\psi_{\vec{n}+\frac{\vec{\mu}-\vec{\mu}'}{2}}$ of action (12) by a gauge link. Because each of the d components of $(\vec{\mu}-\vec{\mu}')/2$ can be equal to ±1 or 0, this gauge link can be constituted by a product $\Pi\, U$ of up to d concatenated elementary orthogonal elementary links. Then there are different paths which connect $\bar{\psi}_{\vec{n}}$ and $\psi_{\vec{n}'}$, and it is convenient to use the average of all the possible paths, or to choose a single path in a random way. With the links included, the I and ∂ terms in eq. (13) are gauge invariant an so is the new action (17).

Including the gauge coupling, the fermion part of the action (17) is extended to,

$$S = a^d \sum_{\vec{n},\vec{n}'} m\bar{\psi}_{\vec{n}} D_{\vec{n},\vec{n}'} e^{i\vec{B}_{n,n'}\cdot(\vec{n}'-\vec{n})}\psi_{\vec{n}'} \ . \tag{21}$$

In the continuum limit $\vec{B}$ tends to a vanishing constant. This action (21) is the one that we propose for the study of light fermions.

In the limit of a constant $\vec{B}$ and using the δ relations (8) the action (21) can be diagonalized in the momentum representation,

$$S = a^d \int_{-\pi}^{\pi} \frac{d^d k}{(2\pi)^d} \widetilde{\bar{\psi}}(k) \widetilde{D}(\vec{k}+\vec{B}) \widetilde{\psi}(k) \tag{22}$$

which is the minimal coupling of the gauge fields to the fermions.

Computing either the propagator or the fermion determinant, is crucial to directly measure chiral symmetry breaking. In the lattice, the Green equation for the propagator,

$$\sum_{\vec{n}''}\left(mI_{\vec{n},\vec{n}''}+\sum_j \gamma_j \partial_{j\,\vec{n},\vec{n}''}\right)(\Pi\, U)_{\vec{n},\vec{n}''}G_{\vec{n}'',\vec{n}'} \\ = I_{\vec{n},\vec{n}'}(\Pi\, U)_{\vec{n},\vec{n}'} \ . \qquad (23)$$

is perfectly defined and can be solved numerically with the standard lattice methods [14]. Although I is not the identity matrix, its nonvanishing matrix elements are very close to the diagonal. The fermion determinant $|D\,U|$ is also calculable. As for the pure gauge term, the Wilson action is unchanged in this case.

However the pole in the action, which was introduced to solve the doubler problem, raises a new difficulty [15] when the gauge coupling is included. This can be understood when we extract pertubatively the coupling of the fermion to a gauge field from eq. (22). The same result can be extracted from the vector Ward identity, and we get,

$$\Gamma_j = \frac{\partial}{\partial p_j}\widetilde{D}(\vec{k}) \\ = i\,\gamma_j\left(1+\tan^2\frac{k_j}{2}\right) . \qquad (24)$$

Now the pole in the fermion Lagrangian (18) will introduce a pole in the coupling say of a photon to an electron. Hence the doublers, which were expelled from the free theory, reappear [15] through the gauge independent interaction. This problem is common to the non-local actions, except for the Bietenholz-Wiese perfect action [12].

IV. A SOLUTION TO THE GINSPARG-WILSON RELATION

We can address at the same token [16] the coupling problem [15], and the Ginsparg-Wilson [7] relation,

$$D\gamma_5+\gamma_5 D = 2rD\gamma_5 D, \qquad (25)$$

where r is a constant of the order of the lattice spacing a. In the case where the action complies with the Ginsparg-Wilson relation, Lüscher [8] showed that the lattice has redefined a chiral symmetry, which includes the chiral anomaly. Moreover, in the continuum limit of $a \to 0$, the chiral symmetry and chiral anomaly of conventional field theory could be recovered [9]. Here we will follow the Chiu-Zenkin [16] prescription to solve the Ginsparg-Wilson relation, which has a linear version for the inverse Dirac matrix,

$$D^{-1}\gamma_5+\gamma_5 D^{-1} = 2r\gamma_5 \ . \qquad (26)$$

A solution of this equation is the sum of an homogeneous solution plus the particular one. Any chiral invariant matrix D_c^{-1} is a solution of the homogeneous equation. We will use the solution of eq. (23) with a vanishing fermion mass m. A simple particular solution is $D_p^{-1} = r$. Our solution is then,

$$D^{-1} = D_c^{-1}+D_p^{-1} \ , \\ D = \frac{D_c}{1+rD_c} \\ = \frac{1}{I+r\sum_j\gamma_j\partial_j}\sum_j \gamma_j\partial_j \ . \qquad (27)$$

In this matrix D, the poles are factorized, this is a simple extension of eq.(23). If we turn off the gauge fields, the Fourier transform can be computed,

$$\widetilde{D} = \frac{i\frac{2}{a}\sum_j\gamma_j\tan\frac{k_j}{2}}{1+r\ i\frac{2}{a}\sum_j\gamma_j\tan\frac{k_j}{2}} \\ = \frac{\sum_j\ i\,\frac{2}{a}\gamma_j\tan\frac{k_j}{2}+r\frac{4}{a^2}\tan^2\frac{k_j}{2}}{1+r^2\frac{4}{a^2}\sum_j\tan^2\frac{k_j}{2}} \ , \qquad (28)$$

which now has a single root, with no other doubles, and no pole in euclidian space.



FIG. 2. On a 24x24 Brillouin lattice with parameters $a = r = 1$ we respectively show the Dirac scalar component (a), and the $i\gamma_2$ component (b) of the Fourier transform of the free action.

It is simpler when $r = a/2$ and in the case where all components k_j vanish except for k_i. In this case eq. (28) is equal to $\left(i\frac{1}{a}\gamma_i \sin k_i + \frac{1}{r}\sin^2\frac{k_i}{2}\right)$, which matches the Wilson action. See Fig. 2 for a plot of the free action in momentum space.

In the case where the fermion has a mass m from the onset, a natural choice for the action consists in using the Dirac matrix D of eq. (27) where we replace the homogeneous solution by the full solution of eq. (23), including the mass term. The action is then,

$$D = \frac{1}{I + r\ \sum_j \gamma_j \partial_j + r\ Im} \left(\sum_j \gamma_j \partial_j + Im \right) \qquad (29)$$

and in the free fermion case the Fourier transform is,

$$\widetilde{D} = \frac{i\frac{2}{a}\sum_j \gamma_j \tan\frac{k_j}{2} + r\frac{4}{a^2}\sum_j \tan^2\frac{k_j}{2} + m + r\,m^2}{1 + r^2\frac{4}{a^2}\sum_j \tan^2\frac{k_j}{2} + r^2 m^2} \ . \qquad (30)$$

V. COMPARING WITH THE NEUBERGER SOLUTION

We now return to the relevant chiral limit of $m = 0$, and compare the factorized action with the literature. The present action is formally very simple, and it is equivalent to the Zenkin action. It is interesting to compare it with the Neuberger action which is presently the most promising solution of the Ginsparg-Wilson relation.

The solution (27) has two interesting regions in the Brillouin zone. In the center we have the single root. There the chiral breaking term vanishes and $D \simeq i\ \not{p}$. The other interesting region is the whole boundary of the Brillouin zone, where the chiral invariant term vanishes and $D \simeq 1/r$. Thus we have a whole surface where $1 - rD = 0$ and this implies that the Chiu-Zenkin condition [16], $det(1 - rD) = 0$ is verified. This is a necessary condition [16] for the topological [8] index

$$tr\gamma_5 D = \text{index } D = n_- - n_+ \qquad (31)$$

to be nonvanishing. The Neuberger [6] action $D_{\mathcal{N}}$ has the same single root, and has $2^d - 1$ points [9] (at the traditional position of the doublers) where $D_{\mathcal{N}} = 1/r$.

It is also straightforward to study the locality of the action D. In the literature we can find different definitions of a local action. Nielsen and Ninomiya called local an action with continuous first derivative in momentum space. The present action D falls in this class. Lüscher is more restrictive [17], and calls local an action of class C^∞, in this class the interactions in position decrease at least exponentially with distance. The Neuberger action belongs to this class. The present action does not, except for dimension $d = 1$, because at the usual position of the doubles its second derivatives do not exist. In $d = 2$ dimensions we Fourier transform back to the position space the free action $\widetilde{D}(k)$, and draw

$$|D|_{\vec{n},\vec{n}'} = |D_0 + i\sum_j \gamma_j D_j|_{\vec{n},\vec{n}'} = \sqrt{\sum_{j=0}^{d} \left| D_{j\,\vec{n},\vec{n}'} \right|^2} \qquad (32)$$

in a logarithmic plot as a function of $|\vec{n} - \vec{n}'|$, see Fig. 3. We find that in the limit of large distances $|D|$ is approximately equal to $exp(-2|\vec{n} - \vec{n}'|^{0.3})$, so the interactions decrease faster than a rational function. The finite range interactions, like the Wilson action, are usually called ultralocal [18], but there are no solutions of the Gisnparg-Wilson relation with an ultralocal rD .



FIG. 3. On a 500x500 lattice with parameters $a = r = 1$ and in the case of free fermions we show $ln(|D|_{\vec{n}-\vec{n}'})$ as a function of $\vec{n} - \vec{n}'$.

The index of our D has been recently computed by Chiu [16,19], who found that D is topologically trivial. The action of Neuberger $D_{\mathcal{N}}$ complies with the index relation. Thus it seems that the action $D_{\mathcal{N}}$ is the best candidate to simulate continuum QCD, including the axial anomaly. Because D has apparently no chiral anomaly, it would be interesting to apply it to the study of chiral symmetry breaking on the lattice with a single flavor.

In continuum QCD the Attiyah-Singer index theorem is verified, but in the lattice this is still an open problem. According to Niedermayer [21], this is scale dependent. If the topological object is larger than the scale of the action then the index theorem should be complied with, however if the topological object has a smaller scale, the index relation is not verified. In Fig. 3 we see that a gauge configuration with finite topological should extend itself for more than 20 or 40 lattice spacings, in order to be larger than the scale of the interaction. In the case of $D_{\mathcal{N}}$, the scale of the fermionic matrix is just a few lattice spacings, an it was checked that $D_{\mathcal{N}}$ complies with the index relation even in a small 6x6 lattice [20]. So it would be interesting to use very large gauge topological objects to check if they would produce a finite index for the fermionic matrix D.

VI. CONCLUSION

With the aim to simulate euclidian QCD, we find a lattice action where the extra roots corresponding to doubled fermions are factorized from the fermion propagator and from the fermion determinant. Like the Wilson fermion action [1], extra hopping terms are included in a initial action. The gauge invariant action with a conventional chiral symmetry has undesired poles in the coupling of a gauge boson to the fermion, but the action with the Ginsparg-Wilson-Lüscher [7,8] symmetry does not suffer from the same pathology. The factorized fermionic action D turns out to be equivalent to the Zenkin [13,16] action. This action complies with the Ginsparg-Wilson [7] relation and with the Chiu-Zenkin [16] criterium. However it is only weakly local and it has been checked that it is topologically trivial, at least in small lattices. The study of this action might help to understand the index relation on the lattice. We conclude that the interest in the present action is just theoretical because the Neuberger action [6] is a better candidate to the simulation of QCD on the lattice.

ACKNOWLEDGMENTS

I thank Michael Creutz, Jean-François Lagae, Jeffrey Mandula and Hugh Shananan for introducing me to the lattice fermion problem during the 1996 Como II conference, and João Seixas and Victor Vieira from CFIF for discussions on lattice techniques. I am indebted to Michael Creutz and to Micha Polikarpov for their remarks and suggestions. This paper was improved by the comment of Wolfgang Bietenholz who pointed out the problems and solutions of non-local lattice actions, and by the comment of Ting-Wai Chiu who called my attention the action of Sergei Zenkin.

APPENDIX A: OTHER POSSIBLE SOLUTIONS WITH DIFFERENT LATTICES

The factorization of doubles can also be applied to different lattices, where the fermion and antifermion fields are placed in different sites. In these cases, unlike the solution in Section II, interpolations are unavoidable in the action, and we no longer have to comply with the Nielsen and Ninomiya theorem. It is possible to find actions without doubles and without the gauge coupling pathology. The gauge links are chosen to directly connect the antifermion sites with the nearest neighbor fermion sites, it is not necessary to concatenate a few links in order to reach a fermion from an antifermion.

An example of a lattice where the same factorization procedure works is a alternate lattice where the ψ and $\bar\psi$ fields are placed at alternate sites. The link structure is simple, however the gauge lattice is 2^d bigger than the lattice of section III.

We will study in more detail another example, where the anti fermion fields $\bar\psi_{\vec n}$ are placed in the normal lattice and the fermion fields $\psi_{\vec n+\vec\mu/2}$ are placed in the dual lattice (or vice versa). For $d>1$, the lattice is composed by plaquettes with 4 elementary links and the Wilson pure gauge action can be used here. However the number of links per zone, the number of links which meet in a vertex and the link length differ from the case of Section III. In the case of Section III these numbers were respectively d, $2d$, and a , while in the present case these numbers are respectively 2^d, 2^d and $a\sqrt{d}/2$. In the case of $d=4$ the number of links is now increased by a factor of 4 and the number of links which meet at a vertex is now doubled. Applying the factorization procedure to the free fermion action,

$$S=\left(\frac{a}{2}\right)^d\sum_{\vec n,\vec\mu}\bar\psi_{\vec n}\Big(m+i\frac{2}{a}\sum_j\mu_j\,\gamma_j\Big)\psi_{\vec n+\frac{\vec\mu}{2}} \tag{A1}$$

$$\widetilde\partial_j(\vec k)=i\frac{2}{a}\tan\frac{k_j}{2}\widetilde I(\vec k)\ ,\quad \widetilde I(\vec k)=\prod_l\cos\frac{k_l}{2}\ ,$$

we find the same momentum representation (28) in the final action.

---

[1] K.G. Wilson, Phys. Rev. **D10** 2445 (1974); *Quarks and Strings on a Lattice*, in New Phenomena in Subnuclear Physics, Erice lectures, July 11-31, 1975, ed. A. Zichichi (plenum, New York, 1977).

[2] J. Kogut and L. Susskind, Phys. Rev. D **11**, 395 (1975).

[3] H. Nielsen and M. Ninomiya, Nuc. Phys. B **185**, 20 (1981); Nuc. Phys. B**193**, 173 (1981).

[4] D. Kaplan, Phys. Lett. B**288**, 342 (1992).

[5] G. 't Hooft, Phys. Lett. **B349**, 491 (1995), hep-th/9411228.

[6] H. Neuberger, Phys. Lett. **B417**, 141 (1998), hep-lat/9707022; Phys. Lett. **B427**, 353 (1998), hep-lat/9801031.

[7] P.H. Ginsparg and K.G. Wilson, Phys. Rev. **D25**, 2649 (1982).

[8] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys. Lett. **B427**, 125 (1998) hep-lat/9801021; M. Lüscher, Phys. Lett. **B428**, 342 (1998) hep-lat/9802011.

[9] K. Fujikawa, Nucl. Phys. **B546**, 480 (1999) hep-th/9811235.

[10] S.D. Drell, M. Weinstein and S. Yankielowicz, Phys. Rev. **D14**, 1627 (1976).

[11] C. Rebbi, Phys. Lett. **186B**, 200 (1987).

[12] W. Bietenholz and U.-J. Wiese, Phys. Lett. **B378**, 222 (1996); hep-lat/9503022. W. Bietenholz and U.-J. Wiese, Nucl. Phys. **B464**, 319 (1996) hep-lat/9510026.

[13] S.V. Zenkin, Bull. Lebedev Phys. Inst. (1988) NO.910; S.V. Zenkin, Mod. Phys. Lett. **A6**, 151 (1991);

[14] J.B. Kogut, Rev. Mod. Phys. **55** (1983) 775.

[15] G.T. Bodwin and E.V. Kovacs, Phys. Lett. **193B**, 283 (1987); G.T. Bodwin and E.V. Kovacs, Phys. Rev. **D35**, 3198 (1987); M. Campostrini, G. Curci and A. Pelissetto, Phys. Lett. **193B**, 279 (1987); A. Pelissetto, Ann. Phys. **182**, 177 (1988);

[16] T. Chiu, Phys. Rev. **D58**, 074511 (1998) hep-lat/9804016; T. Chiu and S.V. Zenkin, Phys. Rev. **D59**, 074501 (1999) hep-lat/9806019; T. Chiu, C. Wang and S.V. Zenkin, Phys. Lett. **B438**, 321 (1998) hep-lat/9806031.

[17] P. Hernandez, K. Jansen and M. Luscher, Nucl. Phys. **B552**, 363 (1999) [hep-lat/9808010].

[18] I. Horvath, Phys. Rev. Lett. **81**, 4063 (1998) [hep-lat/9808002].

[19] T. Chiu, private communication.

[20] T. Chiu, Phys. Rev. **D58**, 074511 (1998) [hep-lat/9804016].

[21] F. Niedermayer, Nucl. Phys. Proc. Suppl. **73**, 105 (1999) hep-lat/9810026.